

ORAL CHEMOTHERAPY EDUCATION



PASSION FOR PATIENTS
NCODA.ORG



ANASTROZOLE

Name of your medication

Generic name — anastrozole (an AS-troh-zole)

Brand name — Arimidex® (uh-RIH-mih-dex)

Common uses

Anastrozole is used in postmenopausal women as:

- A first-step treatment for breast cancer that has spread beyond the breast itself and is either hormone-dependent or has an unknown hormone status
- Treatment for early-stage hormone-dependent breast cancer after surgery with or without radiation therapy
- Treatment for advanced stage breast cancer that has progressed after treatment with tamoxifen (Nolvadex®)

Anastrozole may also be used for other treatments.

Dose and schedule

Taking anastrozole as instructed is important to allow your treatment to be as effective as possible, so here are some key points to remember.

- Your dose may vary, but the usual dose of anastrozole is 1 milligram (1 mg) to be taken by mouth at a scheduled time once a day.
- Anastrozole can be taken with or without food but at the same time each day.
- Anastrozole should be taken whole and not crushed, cut, or dissolved. If you are unable to swallow anastrozole, talk to your care provider or pharmacist for possible options.
- If you miss a dose of anastrozole, **do not** take an extra dose or two doses at one time. Simply take your next dose at the regularly scheduled time. Be sure to write down if you miss a dose and let your care provider know about any missed doses.

Drug and food interactions

- Anastrozole has many drug interactions. Inform your care providers of all prescription medications, over-the-counter medications, vitamins, and herbal products that you are taking.
- Talk with your care team or pharmacist before taking new medications or supplements, or receiving any vaccines.

ORAL CHEMOTHERAPY EDUCATION



PASSION FOR PATIENTS
NCODA.ORG



ANASTROZOLE

Storage and handling

Handle anastrozole with care. Just like when chemotherapy is given into the vein, this drug can be toxic, and exposure of the drug to others should be limited.

- Store anastrozole at room temperature (68°F–77°F) in a dry location away from light.
- Keep anastrozole out of reach of children and pets.
- Leave anastrozole in the provided packaging until it is ready to be taken.
- Whenever possible, give anastrozole to yourself and follow the steps below. If a family member, friend, or caregiver needs to give anastrozole to you, they also need to follow these steps:
 1. Wash hands with soap and water.
 2. Put on gloves to avoid touching the medication. (Gloves are not necessary if you give the drug to yourself.)
 3. Gently transfer the anastrozole from its package to a small medicine or other disposable cup.
 4. Administer the medicine immediately by mouth with water.
 5. Remove gloves and do not use them for anything else.
 6. Throw gloves and medicine cup in household trash.
 7. Wash hands with soap and water.
- If a daily pill box or pill reminder will be used, contact your care team before using:
 - The person filling the box or reminder should wear gloves (gloves are not necessary if you are filling the box or reminder).
 - When empty, the box or reminder should be washed with soap and water before refilling. Be sure to wash hands with soap and water after the task is complete, whether or not gloves are worn
- If you have any unused anastrozole, **do not** throw it in the trash and **do not** flush it down the sink or toilet. Talk to your care provider or pharmacist about proper disposal of anastrozole.
- If you are traveling, put your anastrozole's packaging in a sealed plastic bag. Ask your pharmacist if any additional travel precautions are needed.

ORAL CHEMOTHERAPY EDUCATION



PASSION FOR PATIENTS
NCODA.ORG



ANASTROZOLE

Side Effects of Anastrozole

Below are common side effects that have been known to happen in about one third or more of patients taking anastrozole are listed on the left side of this table. You **MAY NOT** experience these side effects. Options to help manage any side effects that do occur are included on the right side of this table. These should be discussed with your care provider. If you experience any side effect you cannot manage or that is not listed here, contact your care provider.

Possible Side Effect	Management
Hot flashes (sudden sweating and feelings of warmth)	<p>If hot flashes are bothersome:</p> <ul style="list-style-type: none">• Mild, regular exercise may help. Consult first with your care provider.• Try staying in a cool environment.• Wear layers so that if you experience a hot flash, the outer layers may be removed.• Avoid or limit caffeine, spicy food, alcohol, and stress, which may cause hot flashes to worsen.• Inform your care provider if this side effect is bothersome as there may be medications that can help with hot flashes.
Muscle or joint pain or weakness	<ul style="list-style-type: none">• Keep a diary of your pain, including a description of when and where the pain is occurring, what it feels like, and how long it lasts.• Stay as active as possible, but know that it is okay to rest as needed, too.• Tell your care provider if pain interferes with your activity. <p>If the pain or weakness bothers you, ask your provider what you may use to help with this discomfort. Take only pain medication that has been prescribed or recommended by your care provider.</p>

Serious side effects of anastrozole

- Anastrozole may decrease your bone density making you more likely to have a fracture. Your care provider may prescribe medication to help prevent fractures.
- Anastrozole may cause your cholesterol levels to increase in your blood. Speak to your care provider to know when you need to have laboratory tests done to monitor your cholesterol levels. Your care provider may write you a prescription for a cholesterol lowering medication if your levels become high.
- Seek medical attention right away if you have any chest pain or tightness.
- If you have a history of ischemic heart disease, anastrozole may increase your risk of a cardiovascular event.

If you experience ANY uncontrolled side effect, call your physician or healthcare center immediately:

(INSTITUTIONAL CONTACT INFO)

ORAL CHEMOTHERAPY EDUCATION



PASSION FOR PATIENTS
NCODA.ORG



ANASTROZOLE

Handling body fluids and waste

Anastrozole remains in your body for several days after it is taken, so some of the drug may be present in urine, stool, sweat, or vomit. Once you have started to take anastrozole, it is important to follow the instructions below every day for as long as your treatment lasts. This will keep yourself, loved ones, and the environment as safe as possible.

- Pregnant women should avoid touching anything that may be soiled with body fluids from the patient.
- Toilet and septic systems
 - You may use the same toilet, septic tank, and/or sewer that you usually use. If you have a low-flow toilet, close the lid and flush twice to ensure all waste has been discarded.
 - If the toilet or toilet seat becomes soiled with urine, stool, or vomit, clean the surfaces before other people use the toilet.
 - Wash hands with soap and water after using the toilet.
- If you need a bedpan, be sure your caregiver knows to wear gloves to assist with cleanup and to wash the bedpan with soap and water every day.
- If you do not have good control of bladder or bowels, use a disposable pad with a plastic back, a diaper, or a sheet to absorb body waste.
- Wash any skin that has been exposed to body waste or anastrozole with soap and water.
- Linens or clothing that are soiled with body fluids or body waste should be washed separately from other linens and clothing. If you do not have a washer, place the soiled linens in a plastic bag until they can be washed.
- Wash hands with soap and water after touching linens or clothing that may be soiled with body fluids.

Pregnancy, sexual activity, and contraception

- Women should not become pregnant and men should not get a partner pregnant while taking anastrozole. Men and women of childbearing age and potential should use effective contraception during therapy and for a minimum of 3 weeks after the last dose of anastrozole.
- Effective contraception could include 1 or more of the following: oral contraceptive, barrier methods, etc.
- Inform your care provider if you become pregnant.
- Do not breastfeed while taking anastrozole and for 2 weeks after the last dose of anastrozole.
- It is safe to hug and kiss. Special precautions may be needed for sexual activity while on anastrozole, and you are encouraged to ask your care provider.

Obtaining medication

Talk with your care provider about how to obtain your anastrozole.

_____ (PHARMACY OR SPECIALTY PHARMACY CONTACT INFO)

ORAL CHEMOTHERAPY EDUCATION



PASSION FOR PATIENTS
NCODA.ORG



ANASTROZOLE

Additional resources

Anastrozole drug information: www.arimidex.com

Product prescribing information: https://www.accessdata.fda.gov/drugsatfda_docs/label/2011/020541s026lbl.pdf

Product resources: www.arimidex.com/support-breast-cancer.html

Updated – April 12, 2023

Additional instructions

Important notice: The Association of Community Cancer Centers (ACCC), Hematology/Oncology Pharmacy Association (HOPA), National Community Oncology Dispensing Association, Inc. (NCODA), and Oncology Nursing Society (ONS) have collaborated in gathering information for and developing this patient education guide. This guide represents a brief summary of the medication derived from information provided by the drug manufacturer and other resources. This guide does not cover all existing information related to the possible uses, directions, doses, precautions, warnings, interactions, adverse effects, or risks associated with this medication and should not substitute for the advice of a qualified healthcare professional. Provision of this guide is for informational purposes only and does not constitute or imply endorsement, recommendation, or favoring of this medication by ACCC, HOPA, NCODA, or ONS, who assume no liability for and cannot ensure the accuracy of the information presented. The collaborators are not making any representations with respect to the medications whatsoever, and any and all decisions, with respect to such medications, are at the sole risk of the individual consuming the medication. All decisions related to taking this medication should be made with the guidance and under the direction of a qualified healthcare professional.

Permission: Oral Chemotherapy Education (OCE) sheets are provided as a free educational resource for patients with cancer in need of concise, easy-to-understand information about oral cancer drugs. Healthcare providers are permitted to copy and distribute the sheets to patients as well as direct patients to the OCE website for information. However, commercial reproduction or reuse, as well as rebranding or reposting of any type, are strictly prohibited without permission of the copyright holder. Please email permission requests and licensing inquiries to Contact@NCODA.org.

Copyright © 2023 by Hematology/Oncology Pharmacy Association. All rights reserved.